



Depression and Bipolar
Support Alliance

Peer Leadership Center

Fact Sheet

Mental Health Conditions and Mood Disorders

Prevalence

One in four adults – almost 58 million Americans age 18 and older – are diagnosed with a mental health disorder in a given year. Bipolar disorder affects 5.7 million adults, or 2.6 percent, of the U.S. population. Major depressive disorder affects approximately 14.8 million American adults, or about 6.7 percent of the U.S. population age 18 and older in a given year.

Stigma

The vast majority of media stories on mental illness either focus on negative characteristics related to people with the disorder (e.g., unpredictability and unsociability) or on medical treatments. Notably absent are positive stories that highlight recovery of many persons with even the most serious of mental illnesses.

According to *Americans' Views of Mental Health and Illness at Century's End: Continuity and Change, between 1950 and 1996*, "the proportion of Americans who describe mental illness in terms consistent with violent or dangerous behavior nearly doubled."

Americans are hesitant to interact with people who have mental illnesses: 38 percent are unwilling to be friends with someone having mental health difficulties, 64 percent do not want someone who has schizophrenia as a close coworker, and more than 68 percent are unwilling to have someone with depression marry into their family.

Violence

Research has shown that the vast majority of people who are violent do not suffer from mental illnesses.

Mental illness increases the risk of being a victim of murder nearly fivefold. People with severe mental illnesses, schizophrenia, bipolar disorder or psychosis, are 2 ½ times more likely to be attacked, raped or mugged than the general population.

Treatment

Only a third of people with a mental health diagnosis receive treatment.¹


Access to mental health professionals is worse than for other types of doctors. 89.3 million Americans live in a federally-designated Mental Health Professional Shortage Area, and the 2010 National Ambulatory Medical Care survey shows only 2.36% of primary care physician office visits included screening for depression.

In 2010, the United States spent \$135 billion on mental health treatment, or about 5.6 percent of the national health-care spending. Unlike overall health spending, the vast majority of behavioral health services are publicly funded. Medicaid, currently the largest source of financing for behavioral health services in the nation, covers a quarter of all expenditures.

For adults who have health insurance, significant out-of-pocket costs for mental health treatment have risen significantly from 2005 -2009. In a national survey 45% of the untreated cited cost as a barrier.



Figure 2. Out-of-Pocket Costs for Mental Health Services among Adults Aged 18 or Older Who Received Outpatient* or Inpatient** Mental Health Services in the Past Year: 2005 to 2009



Economic Costs

According to the 2004 World Health Report, major depressive disorder is the leading cause of disability in the United States in ages between 15-44. The loss of productivity from mental health disorders costs \$63 billion dollars every year. U.S. workers suffering from bipolar disorder missed double the number of workdays, compared to workers with depression, resulting in disproportionately higher costs to businesses.

The unemployment rate for adults living with mental illness is three to five times higher than for those without mental illness. An estimated \$25 billion is spent annually for disability payments to people with mental illness.

The diagnosis of bipolar disorder in children under 18 years of age rose 4000% from 1996-2006. Over 50% of students with a mental disorder, age 14 and older, drop out of high school—the highest dropout rate of any disability group. High school dropouts cost the U.S. economy \$1.8 billion every year.

Human Costs

Over the past 20 years, dollars spent on inpatient treatment have shifted to outpatient and prescription drugs, with the sickest patients turning up in jails and homeless shelters at a frequency that mirrors that of the late 1800s.

Depressive disorders often co-occur with substance abuse.¹ People living with bipolar disorder have the highest rate of substance abuse of all mental health diagnosis with a rate of 40 to 60 percent. Of adults using homeless services, 31% reported having a combination of these conditions.

More than half of all prison and jail inmates have a mental health problem. 70% of youth in juvenile justice systems have at least one mental disorder, with at least 20% experiencing significant functional impairment from a serious mental illness.

Suicide

Each year more than 8 million Americans think seriously about suicide, more than 1 million attempt suicide and about 38,000 die from suicide. More than 90% of those who die from suicide have a diagnosable mental disorder. 25% to 50% percent of individuals with bipolar disorder attempt suicide at least once during their lifetime; and suicide is a leading cause of death in this group.



Depression and Bipolar
Support Alliance

Peer Leadership Center

References

- ¹ “NIMH: The numbers count—Mental disorders in America.” National Institute of Health.
- ² Substance Abuse and Mental Health Services Administration, (2006) “Violence and Mental Illness: The Facts”
- ³ Americans’ View of Mental Health and Illness at Century’s End: Continuity and Change” Bloomington: Indiana Consortium for Mental Health Services Research and Joseph P. Mailman School of Public Health, Columbia University, 2000.
- ⁴ Pescosolido, B (2007) “Public Perceptions of Mental Illness in 1950 and 1996: What is Mental Illness and Is It to be Feared?” *Journal of Health and Social Behavior* Vol41 188-207
- ⁵ American Psychiatric Association. (1994). *Fact Sheet: Violence and Mental Illness.*” Washington, DC: American Psychiatric Association
- ⁶ Crump C. Mental Disorders and vulnerability to homicidal death: Swedish Nationwide Cohort Study. *BMJ* 2013;346:f55
- ⁷ U.S. Department of Human Services, *Mental Health: A Report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health, 1999
- ⁸ U.S. Department of Human Services, *Designated Mental Health Care Health Professional Shortage Areas*, November 2011.
- ⁹ SAMSHA Spending estimates project, 2010.
- ¹⁰ The Henry J. Kaiser Family Foundation, *Mental Health Financing in the United States: A primer*, April 2011.
- ¹¹ 2004 World Health Report Annex Table 3. Geneva: World Health Organization.
- ¹² U.S. Department of Health and Human Services. *Mental Health: A Report of the Surgeon General*. Rockville, Md., U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, 1999, pp. 408409, 411.
- ¹³ 2006, American Journal of Psychiatry, Dr. Ronald Kessler, Dr. Phillip Wang, Harvard University.
- ¹⁴ Burke-Miller, J.K., *et al.* (April, 2006), “Demographic Characteristics and Employment Among People with Severe Mental Illness in a Multisite Study,” *Community Mental Health Journal* 42(2)
- ¹⁵ Moreno C, Laje G, Blanco C, et al. National trends in the outpatient diagnosis and treatment of bi- polar disorder in youth. *Arch Gen Psychiatry*. 2007;64(9):1032-1039.
- ¹⁶ Alliance for Excellent Education, (2013) "Building a Grad Nation"
- ¹⁷ Interlandi, J, (June 2012), *When My Crazy Father Actually Lost His Mind*, New York Times Magazine.
- ¹⁸ Mcgregor, S. (2007) Substance Abuse and Bipolar Disorder.
- ¹⁹ Burt, M. (2001). “What will it take to end homelessness?” Urban Institute: Washington, D.C., p. 3.
- ²⁰ Glaze, L.& James, D.J. (2006, September). *Mental Health Problems of Prison and Jail Inmates*. US Department of Justice, Office of Justice Programs, Bureau of Justice Statistics: Washington, D.C.
- ²¹ Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web- based Injury Statistics Query and Reporting System (WISQARS) [online]. (2010).
- ²² Risk assessment and psychosocial interventions for suicidal patients. Chesin M, Stanley B., *Bipolar Disorder*. 2013 Jun 20. doi: 10.1111/bdi.12092